

CREATION OF X-RAY HOLES WITH COOL RIMS IN COOLING FLOWS

FABRIZIO BRIGHENTI^{1,2} & WILLIAM G. MATHEWS¹¹University of California Observatories/Lick Observatory, Board of Studies in Astronomy and Astrophysics,
University of California, Santa Cruz, CA 95064
mathews@lick.ucsc.edu²Dipartimento di Astronomia, Università di Bologna, via Ranzani 1, Bologna 40127, Italy
brighenti@bo.astro.it*Draft version February 1, 2008*

ABSTRACT

The density irregularities and holes visible in many *Chandra* X-ray images of cluster and galactic cooling flows can be produced by symmetrically heated gas near the central galactic black hole. As the heated gas rises away from the galactic center, a relatively small number of large plumes and bubbles are formed in qualitative agreement with the observed features. The expanding centrally heated gas drives a shock into the surrounding gas, displacing it radially. Both computational and analytic results show that the ambient gas near the bubble is cooled by expansion, accounting for the cool rims commonly observed around X-ray holes in cooling flows.

Subject headings: galaxies: elliptical and lenticular, CD – galaxies: active – cooling flows – X-rays:
galaxies – galaxies: clusters: general – X-rays: galaxies: clusters

1. INTRODUCTION

High resolution X-ray images of “cooling flows” in elliptical galaxies taken with the *Chandra* observatory indicate that the central hot gas in these systems is not smoothly distributed, but is cavitated on scales comparable to the radio emission (M87: Böhringer et al. 1995; Fabian et al 2001; Wilson et al 2001; M84/NGC 4374: Finoguenov & Jones 2001; NGC 4636: Jones et al. 2001; Loewenstein et al. 2001; David et al. 2002; NGC 507: Forman et al. 2001; NGC 5044: Buote et al. 2002). In view of their short dynamical times, $10^7 - 10^8$ yrs, these disturbances, and by inference also the radio sources, must be highly transient. X-ray holes are also seen in galaxy clusters on scales of $\lesssim 50$ kpc, often approximately coincident with lobes of extended radio emission (e.g. Hydra A/3C295: McNamara et al. 2000; David et al. 2001; Allen et al 2001; Perseus/NGC 1275/3C84: Churazov et al. 2000; Böhringer et al. 1993; Fabian et al 2000; A2052: Blanton et al 2001; A4059: Heinz et al 2002). The Perseus cluster in particular contains a multitude of X-ray holes located at random azimuthal orientations. The X-ray holes are regions in which the gas is either heated or displaced by a non-thermal plasma of comparable energy density. In either case the low-density holes must be buoyant (e.g. Churazov et al. 2001).

One of the most remarkable features of these holes is that the X-ray gas around the rims is often *colder* than the average nearby ambient gas (e.g. Fabian et al. 2000, 2001; Fabian 2002). Low temperature rims have been held as evidence that the surrounding gas has not been (recently) strongly shocked, although shocks must accompany even a subsonic expansion of the hot gas. These low entropy rims cannot be understood as local gas that was shocked and subsequently lost entropy by radiation (Nulsen et al 2002; Soker, Blanton & Sarazin 2002), but may instead be low-entropy gas that has somehow been raised from the center of the flow as these authors suggest.

The random location of older, more distant bubbles in Perseus and the irregular X-ray images within the central few kpcs of elliptical galaxies. are apparently inconsistent with non-thermal jets having fixed orientations defined by the spin axis of massive black holes. Conversely, not every strong radio lobe corresponds to an X-ray hole, as in Perseus/NGC1275/3C84.

We show here that both the random orientation of X-ray holes and their cold rims can be understood if low entropy hot gas near the center of a cooling flow is heated by an active nucleus in the central elliptical galaxy.

2. PREVIOUS STUDIES OF X-RAY HOLES

Several recent theoretical studies of X-ray bubbles have examined the consequences of introducing heated gas in some localized region away from the center of the cooling flow. The 2D hot buoyant bubbles by Churazov et al. (2001), for example, are accompanied by a column of colder (low entropy) gas that moves radially upward near the center of the bubble. These bubbles float upward in the cooling flow atmosphere and come to rest at some large radius where the ambient entropy is equal to that within the bubble. According to Churazov et al, rising bubbles capture cold gas and lift it to large distances from the center, although if this gas is not heated it would be expected to eventually fall back. Nevertheless, the heating required to form the bubbles reduces the global entropy gradient in the cooling flow. Quilis et al. (2001) studied the evolution of a nearly axisymmetric 3D bubble produced by heating gas at some finite radius. They noticed that the buoyant bubble is surrounded by a shell of slightly colder gas which they attributed to cooling expansion as the gas is pushed by the bubble toward regions of decreasing ambient pressure. In these studies gas is assumed to be heated at some off-center site, but the implication is that the ultimate source of energy is located in a central active black hole. Heating by jets that proceed directly from the

AGN core can also create buoyant regions and upwelling of low entropy gas, but only along the jet axis (Reynolds, Heinz, & Begelman 2001; 2002). Reynolds et al. (2002) also note that “curiously, the [observed] cool gas seems to form a shell around the radio lobes and occupies precisely the location where we would expect shock/compressionally heated gas to reside”. Regardless of the heating geometry assumed, by jets or ad hoc off-center bubbles, the global, long-term rate that hot gas cools by radiative losses in cooling flows is unlikely to change without altering the gas density and temperature profiles in disagreement with observations. (Brighenti & Mathews 2002).

3. PRODUCING BUBBLES WITH CENTRAL HEATING

We begin by constructing a simple cooling flow for a massive elliptical galaxy. The gravitational potential is a superposition of a de Vaucouleurs stellar mass profile of total mass $M_{*t} = 7.26 \times 10^{11} M_{\odot}$ and a Navarro-Frenk-White dark halo with virial mass $M_{h,vir} = 4 \times 10^{13} M_{\odot}$ and concentration $c = 10$, typical for the galaxy group environments in which massive E galaxies are thought to form. The initial cooling flow that we consider is generated entirely by mass loss from an old stellar population, $\alpha_* \rho_* = 4.7 \times 10^{-20} (t/t_n)^{-1.3} \text{ gm s}^{-1}$ where $\rho_*(r)$ is the stellar density and $t_n = 13 \text{ Gyrs}$ is the assumed age of the stars. Some heating is provided by Type Ia supernovae, exploding at a rate of $0.06 (t/t_n)^{-1}$ per 100 years per $10^{10} L_B$, each releasing 10^{51} ergs. Supernova heating at this level is consistent with the iron abundance observed in the hot gas but has a relatively minor influence on the overall energetics of the hot gas. The gas-dynamical equations that we solve are identical to those described in Brighenti & Mathews (2002). We use a 2D code modeled after ZEUS (Stone & Norman 1992) with 490 grid zones in both cylindrical coordinates. For $R, z < 50 \text{ kpc}$ the zones are uniform with size $125 \times 125 \text{ pc}$. Outside this region the size increases geometrically toward the outer boundary at $260 \times 260 \text{ kpc}$.

The quiescent cooling flow that results after slowly evolving for 13 Gyrs has a characteristic temperature of $\sim 1 \text{ keV}$ and a gas density profile $\rho \propto r^{-1.2}$ within about 10 kpc and slightly steeper beyond. We assume that gas within $r_h = 1 \text{ kpc}$ of the center is heated at a constant rate $L_h = 2.5 \times 10^{42} \text{ erg s}^{-1}$ for $t_h = 5 \times 10^7 \text{ yrs}$. The heated gas reaches temperatures far in excess of 1 keV and has a sound crossing time much less than t_h , allowing an ultra-hot low-density region to develop near the galactic core. The hot gas pushes outward, driving a shock into the cooling flow gas beyond. The resulting evolution of the gas density and temperature are shown in Figure 1 at three times $t_1 = 40 \times 10^6$, $t_2 = 80 \times 10^6$, and $t_3 = 100 \times 10^6 \text{ yrs}$. The Rayleigh-Taylor (RT) development of a few large plumes filled with heated gas produces irregular X-ray patterns that resemble *Chandra* images of NGC 5044 and other E galaxies. Small wavelength RT unstable features are not prominent even though they would be resolved by the numerical grid. The hot cavity expands gently and is already subsonic ($\lesssim 500 \text{ km s}^{-1}$) at $t = 1 \text{ Myr}$. By time t_1 the weakened precursor shock has moved out to $\sim 20 \text{ kpc}$, moving almost sonically.

Low temperature, bright rims surrounding the hot plumes/bubbles are clearly visible in Figure 1. The peak density in the rims at t_1 is several times that of the ini-

tial unshocked gas at this radius. The minimum entropy $\log(T/n_e^{2/3})$ in the simulation at times $t = 0, t_1, t_2$ and t_3 are 7.15, 7.9, 8.1 and 8.0 respectively. Low entropy gas in the rims at $t \gtrsim t_1$ was originally at $r \sim r_h$ as verified with passively advected tracer particles. This gas cools by adiabatic expansion while rising in the hot gas atmosphere (see also Nulsen et al. 2002; Soker et al. 2002). The larger bubble at time t_2 is penetrated by a disorganized S-shaped column of slowly rising cold gas. The upper panel of Figure 2 shows the X-ray surface brightness of the central cooling flow at time t_2 when the plumes have separated into clearly visible buoyant X-ray bubbles.

It is important to stress that the physical size of the plumes that form is not determined by our computational resolution. Calculations repeated at higher spatial resolution (central grid size=50 pc) also produce a few plumes and (ultimately) bubbles of about the same size as those shown in Figures 1 and 2, which are similar to the X-ray patterns observed. An examination of the effective gravity in the gas rest frame indicates that the outer rims of the large bubbles are RT unstable. At time t_3 the gravitational freefall time across the bubble radius is $t_{ff} \approx 0.1 t_3$; coherent hot bubbles persist for times $\gg t_{ff}$, even at our highest computational resolution.

We performed similar calculations on more extended cooling flows in galaxy groups and clusters with very similar results. For example, the lower panel of Figure 2 illustrates the X-ray surface brightness at time $t = 120 \times 10^6$ years produced in a rich cluster cooling flow ($M_{vir} = 10^{15} M_{\odot}$ and $T_{vir} \approx 6.9 \text{ keV}$) with heating parameters: $L_h = 10^{45} \text{ ergs s}^{-1}$, $r_h = 2 \text{ kpc}$ and $t_h = 5 \times 10^7 \text{ yrs}$. Very little gas cools by radiation losses during the brief duration of our calculations shown in Figures 1 and 2 so the flow is essentially adiabatic. This is expected since a defining attribute of cooling flows is that the dynamical time is much less than the radiative cooling time.

4. SIMILARITY SOLUTION

Cold rims are a generic feature of flows forced to expand into a medium of increasing entropy. This can be illustrated with self-similar flows described by the variable $\eta = r^{-\lambda} t$ (Courant & Friedrichs 1948; Rogers, 1957; Parker, 1961; Chevalier & Imamura 1983). The adiabatic equations for spherical flow can be written in terms of η provided $u = (r/t)U(\eta)$, $\rho = A_o r^{-\alpha} \Omega(\eta)$ and $P = A_o r^{2-\alpha} t^{-2} \Pi(\eta)$. The hot region of centrally heated gas can be represented by an expanding spherical piston at $\eta = \eta_2$ that drives an advancing shock at $\eta = \eta_1 < \eta_2$. The solutions $U(\eta)$, $\Omega(\eta)$ and $\Pi(\eta)$ can be found by inward integration from post-shock values U_1 , Ω_1 and Π_1 which are functions of λ , $\gamma = 5/3$ and the Mach number of the shock $\mathcal{M}$ (e.g. Rogers 1957). We consider the spherical flow that results for pistons uniformly expanding ($\lambda = 1$) into an isothermal, fully ionized medium ($T_o = 10^7 \text{ K}$) with decreasing electron density $n_e = 0.0526 r^{-\alpha} \text{ cm}^{-3}$ where $\alpha = 1.2$ approximates the gas density observed in massive elliptical galaxies (e.g. Brighenti & Mathews 1997). Figure 3 illustrates the flow generated by three pistons driving shocks at $r_1 = 3 \text{ kpc}$ with $\mathcal{M} = 1.5, 2$ and 3. In each case the temperature drops sharply near the piston, *to values even less than T_o* , creating cold rims where the density rises accordingly. Post-shock gas that has just cooled

back to T_o in Figure 3, located at $\log(r/r_1) = -0.179$, -0.108 , and -0.078 for $\mathcal{M} = 1.5$, 2 and 3 respectively, was originally located at $\log(r/r_1) = -0.347$, -0.476 and -0.957 , much closer to the origin. At the piston where $U < 1$ approaches unity, the gas temperature decreases as $\chi \equiv \gamma\Pi/\Omega \propto (1-U)^p$ where $p = 2\alpha/3(5-\alpha) = 0.21$ when $\lambda = 1$. The cold gas adjacent to the piston is a relic of low entropy gas initially near the center that was shocked then cooled by expansion; $d\log\rho/d\log t$ is negative throughout the flowing gas in the plotted solutions. Similarity flows with lower shock Mach numbers have flatter density profiles; for $\mathcal{M} = 1.5$ there is a shallow density minimum at $\log(r/r_1) = -0.047$, indicating weak post-shock RT instability. However, the constant piston velocity assumed in the similarity flows differs from the rapidly decelerating heated bubble in the computed galactic scale flow.

5. FINAL REMARKS AND CONCLUSIONS

If central heating is responsible for the frequently observed X-ray holes and surface brightness fluctuations in typical *Chandra* images, as we claim, then such AGN-black hole heating may be a nearly universal component of cooling flows. The physical origin and nature of this heating – e.g. shock waves, cosmic rays, etc. – has not yet been explored but these details are not essential to the formation of plumes and bubbles. While the production of X-ray irregularities is quite generally insensitive to the assumed heating parameters, we have noticed several trends when L_h , r_h and t_h are varied. Non-linear plumes mature faster when r_h is reduced. If r_h is too large, the cold rims are less pronounced. However, the possibility of learning about such heating details by comparing X-ray images with gasdynamical calculations is limited if the mean time between heating episodes is sufficiently short that the initial pre-heated gas is no longer perfectly quiescent as we have supposed here.

All well-observed cooling flows emit diffuse optical Balmer and forbidden lines visible in the brightest central regions. Enhanced $H\alpha + [\text{NII}]$ emission has been observed near the boundaries of X-ray bubbles (e.g. Blanton et al 2001; Trinchieri & Goudfrooij 2002). It is tempting to conclude that such gas has cooled from the hot phase in the cooler, denser gas around the rims of the plumes or bubbles. But direct cooling from the hot phase is not supported by our detailed hydrodynamic calculations which are essentially unchanged if radiative cooling losses are neglected. The radiative cooling time is greater than the to-

tal time of our simulations. The cooler gas at $T \sim 10^4$ K responsible for the observed optical line emission may simply be displaced by the plumes and bubbles along with the local hot gas. However, on longer timescales than we consider here, heating in the central regions stimulates cooling by thermal instabilities in more distant regions of the flow, so the time-averaged global cooling rate is essentially unchanged (Brighenti & Mathews 2002).

If intermittent AGN heating is reasonably frequent, the entire inner cooling flow will become convective and turbulent (Brighenti & Mathews 2002). Turbulent motions observed in the diffuse optical lines (e.g. Caon et al. 2000) are too energetic on large scales to be understood in terms of supernova explosions, but could result from central heating similar to that described here.

X-ray holes formed by central heating are randomly disposed around the center of the flow, as in the Perseus cluster. This is a particularly desirable feature since the X-ray bubbles need not be aligned with the accretion axis of the central black hole. Nevertheless, radio lobes often appear to be associated with X-ray holes. This connection may be causal, but in some cases the relativistic electrons may readily flow into pre-existing holes; this may be occurring in the central southwestern hole in Perseus (Fig. 1 of Fabian 2001).

Finally, the azimuthally averaged hot gas density profile in these centrally heated models is flattened by the holes, providing an excellent fit to observed X-ray surface brightness profiles which are too centrally peaked otherwise. To maintain this fit, AGN heating would need to reoccur every $\sim 10^8$ years. However, in the presence of central heating the gas temperature profile is less satisfactory, lacking the deep central minimum usually observed.

We conclude that central heating in cooling flows naturally results in a relatively small number of randomly located X-ray bubbles similar to the irregularities observed. In addition, these expanding plumes and bubbles are surrounded by considerably cooler gas, also consistent with recent *Chandra* observations. Gas in the cold rims was shock-heated but subsequently cooled by expansion.

Studies of the evolution of hot gas in elliptical galaxies at UC Santa Cruz are supported by NASA grant NAG 5-8049 and NSF grants AST-9802994 and AST-0098351 for which we are very grateful. FB is supported in part by grants MURST-Cofin 00 and ASI-ARS99-74.

REFERENCES

- Blanton, E. L., Sarazin, C. L., McNamara, B. R. & Wise, M. W. 2001, *ApJ* 558, L15
 Böhringer, H. et al. 1993, *MNRAS*, 264, L25
 Brighenti, F. & Mathews, W. G. 2002, *ApJ* (in press) (astro-ph/0203409)
 Brighenti, F. & Mathews, W. G. 1997, *ApJ* 486, L83
 Buote, D. A. et al. 2002, *ApJ* (in press) (astro-ph/0000000)
 Caon, N., Macchetto, D. & Pastoriza, M. 2000, *ApJ* 553, L125
 Chevalier, R. A. & Imamura, J. N. 1983, *ApJ* 270, 554
 Churazov, E. et al. 2000, *A&A*, 356, 788
 Churazov, E. et al. 2001, *ApJ*, 554, 261
 Courant, R. & Friedrichs, K. O. 1948, *Supersonic Flow and Shock Waves* (Interscience:London), ch. VIC
 David, L. P. et al. 2000, *ApJ*, 557, 546
 David, L., Warmflash, A., Murray, S., & Nulsen, P. E. J. 2002 (*ApJ* in press?) (astro-ph/0108114)
 Fabian, A. C., Sanders, J. S., Ettori, S., Taylor, G. B., Allen, S. W., Crawford, C. S., Iwasawa, K., Johnstone, R. M. & Ogle, P. M. 2000, *MNRAS*, 318, L65
 Fabian, A. C. 2002, (astro-ph/0201386)
 Fabian, A. C. et al. 2000, *MNRAS*, 318, L65
 Fabian, A. C., Sanders, J. S., Ettori, S., Taylor, G. B., Allen, S. W., Crawford, C. S., Iwasawa, K. & Johnstone R. M. 2001, *MNRAS*, 321, L33
 Fabian, A. C., Mushotzky, R. F., Nulsen, P. E. J. & Peterson, J. R. 2001, 321, L20
 Finoguenov, A. & Jones, C. 2001, *ApJ*, 547, L107
 Forman, W., et al. 2001, in *Lighthouses of the Universe Conference*, August 2001 (astro-ph/0111526)
 Heinz, S., Choi, Y.-Y., Reynolds, C.S., Begelman, M.C., 2002, *ApJ*, 569, L79
 Jones, C. et al. 2001, *ApJL* (in press) (astro-ph/0108114)

- Loewenstein, M., Mushotsky, R. F., Angelini, L., Arnaud, K. A. & Quataert, E. 2000, ApJ 555, L21
 McNamara, B. 2000, ApJ, 534, L135
 Nulsen, P. E. J., David, L. P., McNamara, B. R., Jones, C., Forman, W. R. & Wise, M. 2002, ApJ 568, 163
 Parker, E. N. 1961, ApJ 133, 1014
 Quilis, V., Bower, R. G. & Balogh, M. L. 2001, MNRAS, 328, 1091
 Reynolds, C. S., Heinz, S. & Begelman, M. C. 2001, ApJ, 549, L179
 Reynolds, C. S., Heinz, S. & Begelman, M. C. 2002, MNRAS, 332, 271
 Rogers, M. H. 1957, ApJ 125, 478
 Soker, N., Blanton, E. L. & Sarazin C. L. 2002, ApJ (submitted) (astro-ph/0201325)
 Stone, J. M. & Norman, M. L. 1992, ApJS 80, 753
 Trinchieri, G. & Goudfrooij, P. 2002, A&A (in press) (astro-ph/0202416)

FIG. 1.— Evolution of centrally heated gas in a galactic scale cooling flow. The z (vertical) and R (horizontal) coordinates in each panel are labeled in kpc. The upper and lower series of panels show respectively the gas density and temperature at three times: $t_1 = 40 \times 10^6$, $t_2 = 80 \times 10^6$, and $t_3 = 100 \times 10^6$ yrs from left to right. Already at time t_1 the initial precursor shock has moved out to ~ 20 kpc and additional shocks are seen at time t_3 . The hot plumes and bubbles are surrounded by relatively cold gas, particularly at early times. This figure can be viewed in color at the electronic ApJ site.

FIG. 2.— X-ray surface brightness distributions (0.5 - 2 keV) showing bubbles in two flows. The z (vertical) and R (horizontal) coordinates in each panel are labeled in kpc. *Upper panel:* For the galactic scale cooling flow in Figure 1 shown at time $t_2 = 80 \times 10^6$ yrs. *Lower panel:* For a rich cluster cooling flow at time $t = 120 \times 10^6$ yrs with heating parameters: $L_h = 10^{45}$ ergs s^{-1} , $r_h = 2$ kpc and $t_h = 5 \times 10^7$ yrs. This figure can be viewed in color at the electronic ApJ site.

FIG. 3.— Self-similar flow of gas produced by a spherical piston expanding uniformly into an isothermal, fully ionized static gas with initial density $n_e = 0.0526r^{-\alpha} \text{ cm}^{-3}$. In descending order the panels show the radial dependence of the electron density, temperature, pressure and velocity. The flow is shown at the (arbitrary) moment when the shock has arrived at $r_1 = 3$ kpc with Mach numbers 1.5, 2 and 3. The radial variation of the initially undisturbed gas is shown with dashed lines. As suggested by the plots, the temperature in the flowing gas near the piston (at the left of each plot) decreases to zero as the density becomes infinite.





